

Gateway Christian Academy

"A Ministry of Gateway Church"

Certified and Accredited by: ILCS/FLOCS

Bishop Dr. Preston Williams II, Senior Pastor

Shaken Baby Syndrome (Abusive HeadTrauma) Prevention Policy

This policy is designed to prevent the possibility of abusive head trauma during care. Abusive head trauma (also referred to as Shaken Baby Syndrome) occurs in infants and young children, whose neck muscles are not well-developed and whose heads are larger relative to their bodies. As a result, they are especially susceptible to head trauma caused by any type of forceful or sudden shaking, with or without blunt impact.

Damage can occur in as little as 5 seconds.

Abusive head trauma can occur in children up to 5 years of age; however, infants less than one year are at greater risk of injury. Shaken baby syndrome can lead to serious conditions including:

- · Brain damage, problems with memory and attention, cerebral palsy;
- · Blindness or hearing loss;
- · Intellectual, speech or learning disabilities; and
- · Developmental delays.

Signs and Symptoms

The signs and symptoms of shaken baby syndrome or head trauma include:

- · Seizures;
- · Bruises;
- · Lack of appetite, vomiting, or difficulty sucking or swallowing;
- · Lack of smiling or vocalizing;
- · Rigidity, inability to lift the head;
- · Difficulty staying awake, altered consciousness;
- \cdot Difficulty breathing, blue color due to lack of oxygen;
- \cdot Unequal pupil size, inability to focus the eyes or track movement; or
- Irritability.

Injury Prevention

Infant crying is normal behavior, which improves as a child ages. Caregivers should develop proactive strategies to manage stress levels and appropriate responses to a crying child. This includes being selfaware and noticing when the caregiver may become frustrated or angry. Parents/guardians, caregivers and coworkers should discuss what calming strategies are successful with a particular child at home or in the center.

Emergency Response

If a child presents any of the above symptoms or you suspect a baby has suffered abusive head trauma:

- · Call 911, call the parent/guardian and inform your director and management.
- \cdot Report to the appropriate child protective services agency (or law enforcement, if applicable) within

24 hours or less as required by law. See *Child Abuse/ Neglect and Mandated Reporting Policy and Procedure* for further information.

· See *Medical Emergencies-Calling 911* for additional information.

Strategies for Caregivers and Parents

A child is usually shaken out of frustration, often when the child is persistently crying or irritable. The following strategies may work some of the time; but sometimes nothing will comfort a crying child. **A teacher should seek support from a coworker or center management.** If a child is inconsolable on a regular basis, the director and management should be notified and determine if the right supports are in place for the child and for staff.

Do:

 \cdot Hand the child to another caregiver.

[•] Place the child somewhere safe in the classroom (or home) and call the office (or a neighbor) for support; take deep breaths and count to 10.

- \cdot Check to see if the baby's diaper needs changing.
- \cdot Give the baby a bottle. If baby readily takes bottle, feed slowly stopping to burp

often. Do not force the baby to eat.

 \cdot Check for signs of illness and call the parent if you suspect the child is sick.

- · Give baby a pacifier.
- \cdot Hold the baby close against your body and breathe calmly and slowly.
- \cdot Gently rock the baby using slow, rhythmic movements.
- \cdot Sing to the baby or play soft, soothing music.
- \cdot Use "white noise" or rhythmic sounds that mimic the constant whir of noise in the womb
- \cdot Hold the baby on its side or stomach position to help with digestion. Babies $\ensuremath{\textbf{should}}$

always be placed on their backs to sleep.

- \cdot Take the baby for a walk indoors or outside for a ride in the stroller.
- \cdot Be patient: let the baby cry it out if necessary.

Never:

- · Shake a child.
- · Drop a child.
- \cdot Throw a child into the air or into a crib, chair, or car seat.
- \cdot Push a child into any object including walls, doors, and furniture.

Strike a child's head, directly or indirectly.

Resources

In addition to any required state training, the following resources are available to parents/guardians and staff:

Websites:

Abusive Head Trauma-How to Protect Your Baby https://www.healthychildren.org/English/safety-prevention/at-home/Pages/Abusive-Head-Trauma-Shaken-Baby-Syndrome.aspx

National Center on Shaken Baby Syndrome https://www.dontshake.org/

Mary Ann Tocio University (MATU) Resources (can be printed for parents):

Introduction to Early Brain Development https://mybrightweb.brighthorizons.com/EducationPortal/Supplemental %20Materials/ILM Brain%20 Development.pdf Independent Learning Module: Infant Development https://mybrightweb.brighthorizons.com/EducationPortal/Supplemental%20Materials/ILM %20Infant%20 Development.pdf Growing World of Toddlers https://mybrightweb.brighthorizons.com/EducationPortal/Supplemental%20Materials/ ILMToddler%20 Development.pdf Early Brain Development Research Review and Update, Exchange Magazine https://mybrightweb.brighthorizons.com/EducationPortal/Supplemental%20Materials/Early %20Brain%20 Development%20Research%20Review%20and%20Update-Pam%20Schiller.pdf

Related Policies and Procedures

- · Child Abuse/Neglect and Mandated Reporting Policy and Procedure
- · Medical Emergencies-Calling 911
- · Safe Infant and Toddler Handling Procedure

SHAKEN BABY SYNDROME (ABUSIVE HEADTRAUMA) PREVENTION POLICY ACKNOWLEDGEMENT FORM

Parent Signature:_____

Parent Signature: _____

Date:_____

1/30/2022